

(4) LEVEL 7!

OFFICE OF NAVAL RESEARCH

Contract N00014-80-C-0472

Task No. NR 056-749

TECHNICAL REPORT No. 16

On the Synergistic Effects of Laser/Phonon-Stimulated  
Processes: A Master Equation Approach

by

Jui-teng Lin and Thomas F. George

Prepared for Publication

in

Surface Science

University of Rochester  
Department of Chemistry  
Rochester, New York 14627

DTIC  
ELECTE  
DEC 30 1981  
S B D

December, 1981

Reproduction in whole or in part is permitted for any purpose  
of the United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

AD A109025

DTIC FILE COPY

81 12 30 024

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 16	2. GOVT ACCESSION NO. AD A109 005	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) On the Synergistic Effects of Laser/Phonon-Stimulated Processes: A Master Equation Approach		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Jui-teng Lin and Thomas F. George		8. CONTRACT OR GRANT NUMBER(s) N00014-80-C-0472
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Rochester Department of Chemistry Rochester, New York 14627		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 056-749
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program Code 472 Arlington, Virginia 22217		12. REPORT DATE December 1981
		13. NUMBER OF PAGES 12
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of this abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES  Prepared for publication in Surface Science, in press.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) LASER-STIMULATED PROCESSES      ENERGY TRANSFER DYNAMICS ADSPECIES/SURFACE SYSTEM      INTERFERENCE TERMS COHERENT EXCITATION      GENERALIZED MASTER EQUATION SURFACE PHONON EFFECTS      SCHROEDINGER-MARKOFF PICTURE THERMAL INCOHERENT EXCITATION		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The synergistic effects of an adspecies/surface system influenced by a coherent laser field and an incoherent thermal phonon field are theoretically investigated. A generalized master equation is derived in the Schrodinger-Markoff picture and the total stimulated transition rates are decomposed into three parts: thermal phonon, laser and interference terms. The energy transfer dynamics is pictured via the evolution equation of the average excitation of the active mode. A random phase of the off-diagonal matrix elements induced by the laser-stimulated surface processes is introduced. Finally, possible applications of the laser/phonon-stimulated master equation are discussed.		

DD FORM 1 JAN 73 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

On the Synergistic Effects of Laser/Phonon-Stimulated  
Processes: A Master Equation Approach\*

Jui-teng Lin and Thomas F. George<sup>†</sup>  
Department of Chemistry  
University of Rochester  
Rochester, New York 14627 U.S.A.

Abstract

The synergistic effects of an adspecies/surface system influenced by a coherent laser field and an incoherent thermal phonon field are theoretically investigated. A generalized master equation is derived in the Schrödinger-Markoff picture and the total stimulated transition rates are decomposed into three parts: thermal phonon, laser and the interference terms. The energy transfer dynamics is pictured via the evolution equation of the average excitation of the active mode. A random phase of the off-diagonal matrix elements induced by the laser-stimulated surface processes is introduced. Finally, possible applications of the laser/phonon-stimulated master equation are discussed.

Accession For	
NTIS Grant	<input checked="" type="checkbox"/>
NTIS TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
and/or	
Dist	Special
A	

## I. Introduction

Laser-matter interactions of heterogeneous systems have been recently studied and the possible applications/implications of laser-stimulated transient effects, such as materials processing and evaluation, semiconductor lithography and microelectronics and the development of new lasers, motivate the fundamental studies in this newly-developed field.

Laser-stimulated surfaces processes (LSSP) such as migration, desorption, adsorption, ionization, recombination and predissociation usually combine the laser transient effects and the subsequent energy relaxation of the excited adspecies.<sup>1</sup> Due to the multiple time scales of the heterogeneous system, LSSP may be grouped into several types, e.g., mode-selective, adspecies-selective, migrational-selective and local-heating, etc. These selective and nonselective aspects are characterized not only by the coherent properties of the laser radiation, frequency, polarization, intensity and duration, but also by the pumping rate compared to the relaxation or coupling rates among inter- and intra-modes, e.g., electron-phonon interaction, phonon-phonon coupling and active-mode-phonon coupling.<sup>2</sup>

In a previous paper,<sup>1</sup> we studied the desorption dynamics by means of a laser-stimulated rate equation, where the thermal effects and the laser excitation were assumed to be two independent driving forces, and we focused on a cold surface where the laser excitation plays the dominant role. In the present paper, we shall study the synergistic effects of laser/phonon excitation where both the laser and thermal phonons may play equally important roles.

## II. Equation of Motion in the Schrödinger - Markoff Picture

We consider a model system consisting of a group of species adsorbed on a solid surface and subject to laser radiation. The total vibrational Hamiltonian of the system may be expressed in the following form:

$$H = H_A + H_B + H_{AA} + H_{BB} + H_{AB} + H_{AF} + H_{BF}. \quad (1)$$

$H_A$  and  $H_B$  are the unperturbed Hamiltonian of the optically-active (A) mode and the phonon bath (B) modes, respectively;  $H_{AA}$  and  $H_{BB}$  represent the intramode couplings of the active and phonon bath modes, respectively;  $H_{AB}$ ,  $H_{AF}$  and  $H_{BF}$  represent the A and B mode coupling, the A mode and laser field and the B mode and laser field interactions, respectively. For the situation where the vibrational spectrum is partitioned into a high-frequency region consisting of the A modes resonantly excited by a laser, we may neglect  $H_{BF}$  since the low-frequency B modes are far off-resonant with respect to the laser radiation. A second-quantization representation of the total Hamiltonian was previously shown in Ref. 3 consisting of the combinations of the ladder operators  $a^\dagger(a)$  for the active mode and  $b^\dagger(b)$  for the phonon modes. For example,  $H_{AF} = V(t) (a^\dagger + a)$  where  $V(t)$  is proportional to the electric field of the radiation.

In the Schrödinger - Markoff picture (SMP), the equation of motion for the reduced density operator  $S = \text{Tr}_B \rho$ , traced over the phonon modes, with the total Hamiltonian given by eq. (1) is found to be <sup>3,4</sup>

$$\begin{aligned} \frac{\partial S}{\partial t} = & -i[\omega_{\text{eff}} a^\dagger a, S] - iV(t) ([a^\dagger, S] + [a, S]) \\ & + \frac{\gamma_1}{2} (2aSa^\dagger - a^\dagger aS - Sa^\dagger a) + \gamma_1 \bar{n}_j (a^\dagger Sa + aSa^\dagger - a^\dagger aS - Saa^\dagger), \end{aligned} \quad (2.a)$$

with

$$\omega_{\text{eff}}(t) = \omega_0 - \epsilon^*(a^\dagger a + 1) + \phi(t), \quad (2.b)$$

$$\gamma_1 \approx 2\pi \sum_j |K_j|^2 n_j \delta(\omega_0 - 2\epsilon^* n - \sum_j \omega_j), \quad (2.c)$$

$$n_j = \prod_j (\bar{n}_j + 1) - \prod_j \bar{n}_j, \quad (2.d)$$

where  $(\omega_0 - \epsilon^*)$  and  $\omega_j$  are the fundamental frequency of the active mode and the  $j$ -th phonon mode with the vibrational quantum number  $n$  and  $\bar{n}_j$ , respectively. The important features of the above equation of motion in SMP are: (i) the many-body effects of the thermal phonon modes are effectively reduced to a multiphonon damping factor  $\gamma_1$  which is characterized by the mean occupation number  $\bar{n}_j$  (at temperature  $T$ ), the coupling strength  $K_j$  and the delta function  $\delta(\omega_0, \omega_j)$  for conserving energy; (ii) the active mode is treated as a stochastic anharmonic quantum oscillator with an effective frequency  $\omega_{\text{eff}}$  given by an anharmonic correction  $\epsilon^*(a^\dagger a + 1)$  and a random phase or frequency shift  $\phi(t)$ ; <sup>5</sup> (iii) the dephasing effects of the active mode may be caused by a laser/surface-induced mechanism such as thermal fluctuation of the active dipole moment, migration or dislocalization of the adspecies, phase difference between the laser (coherent) and phonon (incoherent) fields coupled to the active mode, random imperfection of the solid surface and finally, the effects of the intramode couplings  $H_{AA}$  and  $H_{BB}$ .

The evolution equation of the density matrix elements for the vibrational states  $(m,n)$ ,<sup>4,6</sup>

$$\frac{\partial S_{mn}}{\partial t} = \frac{1}{i\hbar} \sum_k (H_{mk} S_{kn} - S_{mk} H_{kn}) - \gamma_{mn} (S - \bar{S})_{mn}, \quad (3)$$

can be written for our system, described by the operator equation [eq.(2)], as

$$\begin{aligned} \frac{\partial S_{mn}}{\partial t} = & -i [(\omega_0 - \epsilon^* + \phi)(m-n) - \epsilon^*(m^2 - n^2) - i\gamma_0] S_{mn} \\ & - i \frac{\mu' E(t)}{\hbar} \left[ \sqrt{m} S_{m-1,n} - \sqrt{n+1} S_{m,n+1} + \sqrt{m+1} S_{m+1,n} - \sqrt{n} S_{m,n-1} \right] \\ & + \frac{\gamma_1}{2} \left[ 2 \sqrt{(m+1)(n+1)} S_{m+1,n+1} - (m+n) S_{m,n} \right] \\ & + \gamma_1 \bar{n}_j \left[ \sqrt{mn} S_{m-1,n-1} + \sqrt{(m+1)(n+1)} S_{m+1,n+1} - (m+n+1) S_{m,n} \right], \quad (4) \end{aligned}$$

where  $\mu'$  is a quantity proportional to the derivative of the dipole transition matrix element between ground and first-excited states,  $E(t) = E_0 \cos \omega t$  is the electric field of the linearly polarized laser, and  $\gamma_0 = \frac{1}{2}(\gamma_m + \gamma_n)$  is the mean natural width for the radiative decay from state  $m$  to state  $n$ .

### III. Laser/Phonon-Stimulated Master Equation

The evolution equation of the energy population is given by the diagonal term,  $P_n = S_{nn}$  in eq. (4),

$$\begin{aligned} \frac{dP_n}{dt} = & \gamma_1 \left[ (n+1)(\bar{n}_j+1) P_{n+1} + n\bar{n}_j P_{n-1} - (2n\bar{n}_j + n + \bar{n}_j) P_n \right] - \gamma_n P_n \\ & - (i\mu' E(t)/\hbar) \left[ \sqrt{n} (S_{n-1,n} - S_{n,n-1} - \sqrt{n+1} (S_{n,n+1} - S_{n+1,n})) \right], \quad (5) \end{aligned}$$

where  $P_{n+1} = S_{n+1, n+1}$  and the off-diagonal terms  $S_{n, n+1}$  can be calculated iteratively by means of the equation<sup>6</sup>

$$\frac{\partial S_{mn}^{(j)}}{\partial t} = \frac{1}{i\hbar} [H, S]_{mn}^{(j-1)} - \gamma_{mn} (S - \bar{S})_{mn}^{(j)}, \quad (6)$$

with the condition  $S_{mn}^{(0)}(t) = S_{mn}^{(0)}(t) \delta_{mn}$  and  $S_{mn}^{(0)} = 0$  for  $m \neq n$ , we obtain

$$S_{n, n+1}^{(1)}(t) = -\left(\frac{i}{\hbar}\right) \sqrt{n+1} \int_0^t u' E(t') \exp \left[ \int_0^{t'} \Omega(t'') (t' - t'') dt'' \right] \times \\ [P_{n+1}^{(0)}(t') - P_n^{(0)}(t')] dt', \quad (7.a)$$

where

$$\Omega(t) = i[\omega_0 - 2\varepsilon^*(n+1) + \phi(t)] - (2n+1-4n\bar{n}+4\bar{n})\gamma_1/2 - \gamma_0. \quad (7.b)$$

Note that  $\Omega(t)$  contains a random phase factor  $\phi(t)$ . By assuming a "white noise",  $\langle \phi(t) \rangle = 0$ , and a Markoffian correlation,  $\langle \phi(t) \phi(t') \rangle = 2\gamma_2 \delta(t-t')$ , we may evaluate the ensemble-averaged quantity [over the random variable  $\phi(t)$ ], as

$$\langle \exp \left[ i \int_0^{t'} \phi(t'') (t' - t'') dt'' \right] \rangle \approx \sum_{n=0}^{\infty} \frac{(-\gamma_2 t)^n}{n!} = e^{-\gamma_2 t}, \quad (8)$$

which is governed by the dephasing ( $T_2$ ) factor  $\gamma_2$ . Equation (7.a) then takes the form

$$\langle S_{n, n+1}^{(1)}(t) \rangle = -\left(\frac{i}{\hbar}\right) \sqrt{n+1} \int_0^t u' E(t') \exp[\bar{\Omega}_1] [\langle P_{n+1}^{(0)}(t') \rangle - \langle P_n^{(0)}(t') \rangle] dt', \quad (9.a)$$

$$\bar{\Omega}_1(t-t') = \left( i[\omega_0 - 2\varepsilon^*(n+1)] - \Gamma_1 \right) (t-t') \quad (9.b)$$

$$\Gamma_1 = \gamma_1(2n+1+4n\bar{n}+4\bar{n})/2 + \gamma_2 + \gamma_0. \quad (9.c)$$



By the same procedure, we find

$$\langle S_{n-1,n}^{(1)}(t) \rangle = \left(\frac{i}{\hbar}\right) \sqrt{n} \int_0^t \mu' E(t') \exp[\bar{n}_2] [\langle P_{n-1}^{(0)}(t') \rangle - \langle P_n^{(0)}(t') \rangle] dt', \quad (10.a)$$

$$\bar{n}_2(t-t') = \left( i[\omega_0 - 2\varepsilon^* n] - \Gamma_2 \right) (t-t'), \quad (10.b)$$

$$\Gamma_2 = \gamma_1(2n-1+4n\bar{n})/2 + \gamma_2 + \gamma_0. \quad (10.c)$$

Substituting eqs. (9) and (10) into eq. (5), we obtain the ensemble-averaged generalized maser equation (GME), neglecting the ensemble-averaged notation  $\langle \dots \rangle$ , for the first-order solution,

$$\frac{dP_n}{dt} = \gamma_1 [(n+1)(\bar{n}_j+1)P_{n+1} + n\bar{n}_jP_{n+1} - (2n\bar{n}_j+n+\bar{n}_j)P_n] - \gamma_n P_n - \int_0^t dt' \left( W_{n,n-1}(t-t') [P_{n-1}(t') - P_n(t')] + W_{n,n+1}(t-t') [P_{n+1}(t') - P_n(t')] \right), \quad (11)$$

with the time dependent transition rates

$$W_{n,n+1}(t-t') = 2(n+1)(\mu'/\hbar)^2 E(t)E(t') e^{-\Gamma_1(t-t')} \cos[(\omega_0 - 2\varepsilon^* n - 2\varepsilon^*)(t-t')], \quad (12.a)$$

$$W_{n,n-1}(t-t') = 2n(\mu'/\hbar)^2 E(t)E(t') e^{-\Gamma_2(t-t')} \cos[(\omega_0 - 2\varepsilon^* n)(t-t')]. \quad (12.b)$$

The above GME may be reduced to the Pauli master equation (ME) by using the fact that the population inversion  $[P_{n+1}(t) - P_n(t)]$  is usually a slowly varying function<sup>4</sup> in the time of  $\Gamma_{1,2}^{-1}$  and hence can be factored outside the integral in eq. (11). Taking an average over the cycle of the field,  $2\pi/\omega$ , and assuming a near resonance condition such that the integrated transition rates  $W_{n,n+1}(t)$  become time-independent, and we obtain a simple laser/phonon-stimulated master equation (LPME)

$$\begin{aligned} \frac{dP_n}{dt} = & \left( W_{n,n+1}^P + W_{n,n+1}^{LP} \right) (P_{n+1} - P_n) + \left( W_{n,n-1}^P + W_{n,n-1}^{LP} \right) (P_{n-1} - P_n) \\ & + \left( R_{n,n+1} P_{n+1} - R_{n,n-1} P_n \right) - \gamma_n P_n, \end{aligned} \quad (13)$$

where the stimulated transition rates are given by

$$W_{n,n+1}^P = \gamma_1 \bar{n}_j (n+1), \quad (14.a)$$

$$W_{n,n-1}^P = \gamma_1 \bar{n}_j n, \quad (14.b)$$

$$W_{n,n+1}^{LP} = \frac{(n+1)}{2} \left( \frac{\mu' E_0}{\hbar} \right)^2 g_1(\bar{\Delta}), \quad (14.c)$$

$$W_{n,n-1}^{LP} = \frac{n}{2} \left( \frac{\mu' E_0}{\hbar} \right)^2 g_2(\bar{\Delta}), \quad (14.d)$$

$$g_{1,2}(\bar{\Delta}) = \Gamma_{1,2} / (\bar{\Delta}_{1,2}^2 + \Gamma_{1,2}^2), \quad (14.e)$$

$$\bar{\Delta}_1 = \omega_0 - 2\varepsilon^* (n+1) - \omega, \quad (14.f)$$

$$\bar{\Delta}_2 = \omega_0 - 2\varepsilon^* n - \omega. \quad (14.g)$$

The phonon-induced relaxation rates  $R_{n,n+1} = (n+1)\gamma_1$  and  $R_{n,n-1} = n\gamma_1$  describe the irreversible energy dissipative processes for excited-species on a cold surface. The final term in eq.(13) describes the spontaneous radiative decay of the  $n$ -th vibrational levels.

#### IV. Discussion and Conclusion

It is shown in the LPME [eq.(10)] that the active mode is influenced by a coherent laser field and an incoherent thermal phonon field with total stimulated transition rates  $(W_{n,n+1}^P + W_{n,n+1}^{LP})$  which may be further decomposed into three components  $(W_{n,n+1}^P + W_{n,n+1}^L - W_{n,n+1}^I)$ , corresponding to phonon-stimulated, laser-stimulated and the "interference" transition rates, respectively.  $W_{n,n+1}^P$  is given in eq.(14) and

$$W_{n,n+1}^L = \left( \frac{n+1}{2} \right) \left( \frac{\mu' E_0}{\hbar} \right)^2 \mathcal{L}_1, \quad (15.a)$$

$$W_{n,n-1}^L = \left(\frac{n}{2}\right) \left(\frac{\mu' E_0}{\hbar}\right)^2 \mathcal{L}_2, \quad (15.b)$$

$$\mathcal{L}_{1,2} = \gamma_0 / (\bar{\Delta}_{1,2}^2 + \gamma_0^2), \quad (15.c)$$

are simply the surface-free laser pumping rates characterized by the Rabi frequency and a Lorentzian with FWHM equal to the natural level width  $\gamma_0$ . The interference term is related to  $W^L$  and  $W^P$  by

$$W_{n,n\pm 1}^I \approx \left(W_{n,n\pm 1}^L\right)^2 \left(\gamma_2 + W_{n,n\pm 1}^P N_{1,2} / \bar{n}_j(n+1)\right), \quad (16)$$

for  $\gamma_0 \gg N_{1,2} \gamma_1 + \gamma_2, \bar{\Delta}_{1,2}$ .

For a monatomic lattice with phonon frequency  $\approx$  Debye frequency ( $\omega_D$ ), the multiphonon coupling factor [eq. (2.c)] may be approximated as  $\gamma_1 \approx \gamma_1(0) T^{N-1} [\exp(\hbar\omega_0/kT) - 1]$ , with the zero temperature factor  $\gamma_1(0) \sim \alpha^{2N}$ , where  $N \approx (\omega_0 - 2\varepsilon^*n)/\omega_D$  in the order parameter and  $0 < \alpha \ll 1$ . For a highly excited active mode, the multiphonon order parameter,  $N$ , decreases as the anharmonic correction  $2\varepsilon^*n$  increases and in turn results a stronger coupling factor  $\gamma_1$ . A stronger coupling factor  $\gamma_1$  results in a significant interference term  $W^I$  as shown in eqs. (16). Therefore, the total transition rate in the LPME can be approximated by the sum of  $W^P$  and  $W^L$  only when  $W^I$  is negligible, i.e.,  $(N_{1,2}\gamma_1 + \gamma_2) \ll \gamma_0$ .

To give a clearer picture of the synergistic effects of laser/phonon excitation processes, we consider the average excitation of the active mode,  $\langle n_A \rangle = \langle a^\dagger a \rangle = \langle \text{Tr}(S a^\dagger a) \rangle$ , where  $\langle \dots \rangle$  denotes an ensemble average over the A and B modes. The evolution of  $\langle n_A \rangle$  can be obtained from eq. (2) as

$$\langle n_A \rangle = [i\mu' E_0 t / \hbar] [\langle a \rangle - \langle a^\dagger \rangle] + \gamma_1 \bar{n}_j - \gamma_1 \langle n_A \rangle, \quad (17)$$

which shows that the active mode is pumped by two forces, a laser field  $\mu'E(t)$  and a thermal phonon field  $\gamma_1 \bar{n}_j$ , while it is damped by a multiphonon relaxation rate  $\gamma_1$ , the last term in eq. (17). For the case of no laser field,  $E(t)=0$ ,  $\langle n_A \rangle$  reaches the steady-state value  $\bar{n}_j$  and is in thermal equilibrium with all the other modes. With laser radiation, a nonequilibrium state where the active mode has a higher vibrational energy than the others is provided, and selective excitation is possible in a time scale shorter than that of the multiphonon relaxation  $\gamma_1^{-1}$ .

In conclusion, we propose some possible applications of LPME [eq. (13)] where the laser and the surface play equally important roles: (i) solving the LPME for  $P_n$  by, e.g., generating function method, enables us to calculate physical quantities such as the desorption probability<sup>1</sup> and the desorption rate (or the mean first passage time);<sup>8</sup> (ii) the adspecies/surface spectrum governed by the correlation function of the off-diagonal matrix elements [eqs. (9) and (10)] will provide us surface dynamical information such as the surface coverage and the migration rate of the adspecies through the dephasing broadening of the spectrum  $\gamma_2$  which reflects the collisional effects; (iii) replacing the vibrational population function by the lattice site probability, the LPME in eq. (13) then may be used to describe the dynamical motion of the adspecies on the solid surface in which the vibrational pumping rates may be regarded as the lattice hopping rates.<sup>9</sup>

One of us (J. L.) wishes to thank Prof. J. H. Eberly and Dr. K. S. Lam for useful conversations.

\*This work was supported in part by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Grant AFOSR 82-0046. The United States Government is authorized to reproduce and distribute reprints.

<sup>†</sup>Camille and Henry Dreyfus Teacher-Scholar, 1975-82.

1. J. Lin and T. F. George, Chem. Phys. Lett. 66, 5 (1979).
2. J. Lin and T. F. George, Surface Sci. 100, 381 (1980); 107, 417 (1981) and references therein.
3. J. Lin, A. C. Beri, M. Hutchinson, W. C. Murphy and T. F. George, Phys. Lett. 79A, 233 (1980).
4. W. H. Louisell, Quantum Statistical Properties of Radiation, (Wiley, New York, 1973), Chap. 6.
5. M. Sargent III, M. O. Scully and W. E. Lamb, Jr., Laser Physics (Addison-Wesley, Reading, MA, 1974), Chap. 7.
6. A. Yariv, Quantum Electronics (Wiley, New York, 1975), Appendix 4.
7. A. Nitzan, S. Mukamel and J. Jortner, J. Chem. Phys. 60, 3929 (1974).
8. E. W. Montroll and K. E. Shuler, Adv. Chem. Phys. 1, 361 (1958).
9. J. Lin and T. F. George, J. Phys. Chem. 84, 2957 (1980).

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, N.C. 27709	1
ONR Western Regional Office Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Eastern Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. David L. Nelson Chemistry Program Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, 056

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. G. A. Somorjai Department of Chemistry University of California Berkeley, California 94720	1	Dr. C. P. Flynn Department of Physics University of Illinois Urbana, Illinois 61801	1
Dr. L. N. Jarvis Surface Chemistry Division 4555 Overlook Avenue, S.W. Washington, D.C. 20375	1	Dr. W. Kohn Department of Physics University of California (San Diego) LaJolla, California 92037	1
Dr. J. B. Hudson Materials Division Rensselaer Polytechnic Institute Troy, New York 12181	1	Dr. R. L. Park Director, Center of Materials Research University of Maryland College Park, Maryland 20742	1
Dr. John T. Yates Department of Chemistry University of Pittsburgh Pittsburgh, Pennsylvania 15260	1	Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455	1
Dr. Theodore E. Madey Surface Chemistry Section Department of Commerce National Bureau of Standards Washington, D.C. 20234	1	Dr. Chia-wei Woo Department of Physics Northwestern University Evanston, Illinois 60201	1
Dr. J. M. White Department of Chemistry University of Texas Austin, Texas 78712	1	Dr. D. C. Mattis Polytechnic Institute of New York 333 Jay Street Brooklyn, New York 11201	1
Dr. Keith H. Johnson Department of Metallurgy and Materials Science Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. Robert M. Hexter Department of Chemistry University of Minnesota Minneapolis, Minnesota 55455	1
Dr. J. E. Demuth IBM Corporation Thomas J. Watson Research Center P.O. Box 218 Yorktown Heights, New York 10598	1	Dr. R. P. Van Duyn Chemistry Department Northwestern University Evanston, Illinois 60201	1

# TECHNICAL REPORT DISTRIBUTION LIST, 056

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. S. Sibener Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637	1	Dr. Martin Fleischmann Department of Chemistry Southampton University Southampton SO9 5NH Hampshire, England	1
Dr. M. G. Lagally Department of Metallurgical and Mining Engineering University of Wisconsin Madison, Wisconsin 53706	1	Dr. J. Osteryoung Chemistry Department State University of New York at Buffalo Buffalo, New York 14214	1
Dr. Robert Gomer Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637	1	Dr. G. Rubloff I.B.M. Thomas J. Watson Research Center P. O. Box 218 Yorktown Heights, New York 10598	1
Dr. R. G. Wallis Department of Physics University of California, Irvine Irvine, California 92664	1	Dr. J. A. Gardner Department of Physics Oregon State University Corvallis, Oregon 97331	1
Dr. D. Ramaker Chemistry Department George Washington University Washington, D.C. 20052	1	Dr. G. D. Stein Mechanical Engineering Department Northwestern University Evanston, Illinois 60201	1
Dr. P. Hansma Chemistry Department University of California, Santa Barbara Santa Barbara, California 93106	1	Dr. K. G. Spears Chemistry Department Northwestern University Evanston, Illinois 60201	1
Dr. P. Hendra Chemistry Department Southampton University England SO9JNH	1	Dr. R. W. Plummer University of Pennsylvania Department of Physics Philadelphia, Pennsylvania 19104	1
Professor P. Skell Chemistry Department Pennsylvania State University University Park, Pennsylvania 16802	1	Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 44106	2
Dr. J. C. Hemminger Chemistry Department University of California, Irvine Irvine, California 92717	1	Professor D. Hercules University of Pittsburgh Chemistry Department Pittsburgh, Pennsylvania 15260	1



TECHNICAL REPORT DISTRIBUTION LIST, 056

No.  
Copies

Professor N. Winograd  
The Pennsylvania State University  
Department of Chemistry  
University Park, Pennsylvania 16802 1

Professor J. F. George  
The University of Rochester  
Chemistry Department  
Rochester, New York 14627 1

Professor Dudley R. Herschbach  
Harvard College  
Office for Research Contracts  
1350 Massachusetts Avenue  
Cambridge, Massachusetts 02138 1

Professor Horia Metiu  
University of California,  
Santa Barbara  
Chemistry Department  
Santa Barbara, California 93106 1

Professor A. Steckl  
Rensselaer Polytechnic Institute  
Department of Electrical and  
Systems Engineering  
Integrated Circuits Laboratories  
Troy, New York 12181 1

Professor R. D. Archer  
University of Massachusetts  
Chemistry Department  
Amherst, Massachusetts 01003 1

Dr. A. C. Pastor  
Hughes Research Laboratories  
3011 Malibu Canyon Road  
Malibu, California 90265 1